





GPRA CLINICAL INDICATORS

Santa Ynez Band of Mission Indians
2001, 2002, 2003 GPRA Plus Data


Indicator 2a: The percentage of Santa Ynez Indian diabetic patients who received the HbA1c test, to measure glycemic control, in the prior year.




Indicator 2b: The percentage of Santa Ynez Indian
diabetic patients who received the HbA1c test
and had ideal control (<7) in the prior year.




Indicator 2c: The percentage of Santa Ynez Indian diabetic patients who received the HbA1c test and had poor control (>9.5) in the prior year.




Indicator 3: The percentage of Santa Ynez Indian
diabetic patients who achieved
blood pressure control (<130/80) in the prior year.




Indicator 4: The percentage of Santa Ynez Indian diabetic patients who have been assessed for dyslipidemia (LDL cholesterol) in the prior year.




Indicator 5: The percentage of Santa Ynez Indian
diabetic patients who have been
assessed for nephropathy in the prior year.




Indicator 6: The percentage of Santa Ynez Indian diabetic patients who have been assessed for an annual retinal examination in the prior year.




Indicator 14: The percentage of Santa Ynez Indian diabetic patients with an annual dental visit in the prior year.




Indicator 7: The percentage of Santa Ynez Indian women
21 to 64 years old who had a PAP smear
or refused a PAP smear within the past three years.



Indicator 8: The percentage of Santa Ynez Indian women 52 to 69 years old who received a mammogram or refused a mammogram within past three years.



Indicator 24: The percentage of influenza vaccinations administered to Santa Ynez Indian men and women 65 years old and older within the past year.



Indicator 25: The percentage of pneumococcal vaccinations administered to Santa Ynez Indian men and women age 65 and older any time in the past.

